

A new species of the spider genus *Harpactea* Bristowe, 1939  
from Azerbaijan (Aranei Dysderidae).

Новый вид пауков рода *Harpactea* Bristowe, 1939  
из Азербайджана (Aranei Dysderidae).

P.M. Dunin  
П.М. Дунин

Institute of Ecology of the Volga River Basin, Russian Academy of Sciences, Komzina Street 10, Togliatti – 3, Samara Area 445003 Russia.

Институт экологии Волжского бассейна РАН, ул. Комзина, 10, г.Тольятти – 3 Самарской обл., 445003 Россия.

KEY WORDS: Aranei, Dysderidae, *Harpactea* new species, Azerbaijan.

КЛЮЧЕВЫЕ СЛОВА: Aranei, Dysderidae, новый вид *Harpactea*, Азербайджан.

РЕЗЮМЕ: Описание нового паука из аридных арчево-фисташковых редколесий Большого Кавказа Азербайджана. *Harpactea dashdamirovi* sp.n. принадлежит к подгруппе «caucasia» группы видов «hombergi», очевидно, ближе всего к *H.nachitschevanica* Dunin, 1991, но отличается не треугольным, а овальным бульбусом, слабее закрученным эмболюсом, несколько более коротким и массивным кондуктором, а также неболь-

шими деталями размеров, окраски и вооружения конечностей.

ABSTRACT: A new spider from arid *Juglans-Pistacia* sparse stands of the Caucasus Major of Azerbaijan is described. *Harpactea dashdamirovi* sp.n. belongs to the *caucasia*-subgroup of the *hombergi*-group, being perhaps especially close to *H.nachitschevanica* Dunin, 1991, but differing in




Fig. *Harpactea dashdamirovi*, male palp.

Рис. *Harpactea dashdamirovi*, пальпс самца.

the oval, not triangular, bulbus, less dramatically coiled embolus, somewhat shorter and stouter conductor, as well as by certain minor details of the size, coloration, and leg spinulation.

No sooner had my revision of the Caucasian Dysderidae been issued [Dunin, 1992], with a total of 28 *Harpactea* spp. recognized, I received small material from Azerbaijan containing another new species of that genus. This paper is devoted to its description, entirely following the pattern accepted in Dunin [1992]. Type material is kept in the Zoological Museum of the State University of Moscow, Russia.

*Harpactea dashdamirovi*, sp.n.

Fig.

Holotype: ♂, Azerbaijan, Agdash Distr., Bozdagh Mt. Range, Turianchai Reserve, Dadly-Bulag Valley, 100-300 m a.s.l., arid *Juglans-Pistacia* sparse forest with *Quercus* remains, litter and under stones, 8.V.1992; H. Aliev, S. Dashdamirov & V. Gusarov. - Paratype: ♂, same locality, 6.V.1992; leg. S. Dashdamirov.

Etymology: Honours Dr. Selvin D. Dashdamirov, one of the collectors.

Diagnosis: The new species belongs to the *caucasia*-subgroup of the *hombergi*-group, being perhaps especially close to *H.nachitschevanica* Dunin, 1991, but differing in the oval, not triangular,

bulbus, less strongly coiled embolus, somewhat shorter and stouter conductor, as well as by certain minor details of the size, coloration, and leg spinulation.

Description: Male. Carapace 1.53 long, 1.20 wide, cephalic portion 0.63 wide. Carapace brown, finely shagreened. Sternum dark yellow. Chelicerae brown, with small dark brown spots. Labium and gnathocoxae yellow-brown. Legs yellow to dark yellow. Leg armature: Fm I Ad(2), Fm II Ad(1.1), Fm III Dm(1), ADd(1.1)/(1), Fm IV Db(1.1)/(1.1.1); Pt III A(1); Ti III-IV AD(1.1.1), AV(1.1.1), PD(1.1.1), PV(1.1.1). Abdomen 1.86 long, 1.01 wide. Abdomen grey-yellow, with short hairs. Copulatory organ as in Figs 1-2.

Female unknown.

ACKNOWLEDGEMENTS. I am greatly indebted to my colleague friend Mr. S.D. Dashdamirov (Baku), who has been supplying me with his copious spider collections mainly deriving from the Caucasus. In addition, Dr. S.I. Golovatch (Moscow) kindly checked the English of the final draft. This contribution has partly been sponsored by the Soros Foundation.

## References

- Dunin P.M. 1992. [The spider family Dysderidae of the Caucasian fauna (Arachnida Aranei Haplogynae)] // *Arthropoda Selecta*. Vol.1. No.3. P.35-76 [In Russian, English summary].