

New records of the linyphiid spiders from the Bureinsky Nature Reserve, Khabarovsk Province, Russian Far East (Arachnida: Aranei: Linyphiidae)

Новые находки пауков-линифиид из Буреинского заповедника (Хабаровский край, Дальний Восток России) (Arachnida: Aranei: Linyphiidae)

**L.A. Trilikauskas¹ & A.V. Tanasevitch²
Л.А. Триликаускас¹, А.В. Танасевич²**

¹Bureinsky State Nature Reserve, Zelionaya Str.3, Chegdomyn, Khabarovsk Province 682030 Russia.

Буреинский государственный природный заповедник, ул. Зеленая 3, Чегдомын, Хабаровский край 682030 Россия.

²Centre for Forest Ecology and Productivity, Russian Academy of Sciences, Profsoyuznaya Str. 84/32, Moscow 117997 Russia.

Центр по проблемам экологии и продуктивности лесов РАН, Профсоюзная ул. 84/32, Москва 117997 Россия.

KEY WORDS: Spiders, Linyphiidae, Siberia, Khabarovsk Province, faunistics, distribution.

КЛЮЧЕВЫЕ СЛОВА: Пауки, Linyphiidae, Сибирь, Хабаровский край, фаунистика, хорология.

ABSTRACT. A checklist of 75 linyphiid species collected in the Bureinsky Nature Reserve, Khabarovsk (= Habarovsk) Prov. and adjacent territories is compiled, with 32 species recorded as new to the region and one, *Nerienne oidedicata* van Helsdingen, 1969, as new to the Russian fauna. Habitat data are provided for each species listed. Four previous misidentifications are rectified (correct names on the right): *Bathyphantes simillimus* (L. Koch, 1879) = *Bathyphantes eumenis* (L. Koch, 1879); *Ceratinella brevipes* (Westring, 1851) and *Ceratinella scabrosa* (O. Picard-Cambridge, 1871) = *Ceratinella wideri* (Thorell, 1871), *Nenilinium luteolum* (Loksa, 1965) = *Nenilinium asiaticum* Eskov, 1988. Based on the new records, the linyphiid fauna of the nature reserve and its adjacent parts comprises 143 species.

РЕЗЮМЕ. Приведен список из 75 видов пауков семейства Linyphiidae, найденных в Буреинском заповеднике и его окрестностях. Из них 32 видов отмечены здесь впервые, один вид, *Nerienne oidedicata* van Helsdingen, 1969, — впервые для фауны России. Для всех видов указаны новые места находок. Исправлены четыре допущенные ранее ошибки в определениях (правильные названия справа): *Bathyphantes simillimus* (L. Koch, 1879) = *Bathyphantes eumenis* (L. Koch, 1879); *Ceratinella brevipes* (Westring, 1851) and *Ceratinella scabrosa* (O. Picard-Cambridge, 1871) = *Ceratinella wideri* (Thorell, 1871); *Nenilinium luteolum* (Loksa, 1965) = *Nenilinium asiaticum* Eskov, 1988. С учетом новых данных, фауна пауков семейства Linyphiidae заповедника и его окрестностей насчитывает 143 вид.

Introduction

Recently, a checklist of 104 linyphiid species of the Bureinsky State Nature Reserve and adjacent territories has been published [Tanasevitch & Trilikauskas, 2004]. New material collected in 2004 and 2005, coupled with a restudy of the earlier undetermined samples, allows to essentially update and complete this list, also providing useful habitat information concerning the nature reserve's spiders.

Material and methods

The present paper is based on new or previously undetermined material collected by L. A. Trilikauskas in 1999–2005, by D. V. Logunov in July and August 2004, as well as by A. V. Tanasevitch in May and June 2003 in the Bureinsky State Nature Reserve and its adjacent territories. The material is stored in the Zoological Museum of the Moscow State University, Moscow and the Bureinsky Nature Reserve, Chegdomyn.

The species recorded in the region for the first time are marked with asterisk.

ABBREVIATIONS. The following abbreviations have been used in the text: B.R. — Bureinsky Nature Reserve, Khabarovsk Province; L. — “Levaya” (= left) Bureya River, P. — “Pravaya” (= right) Bureya River.

List of linyphiid spiders

Agyneta affinisoides Tanasevitch, 1984*

MATERIAL. 1 ♂, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. Middle and East Siberia, Cisokhotia; through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk Prov. Known also from Mongolia.

Agyneta conigera (O. Pickard-Cambridge, 1863)*

MATERIAL. 2 ♂♂, B.R., Bureya River Valley, 1 km downstream of confluence of P. and L. Bureya River, 21–26.VI.2004, 1–6.VII.2004, leg. L. Trilikauskas; 1 ♂, B.R., P. Bureya River Valley, 300 m upstream of the mouth, 2–7.VII.2004, leg. L. Trilikauskas.

DISTRIBUTION. Europe, including the tundra zone, the Caucasus; in Siberia: from the Urals eastward to East Siberia, Cisokhotia. Eastern Kazakhstan, through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk Prov. Known also from Mongolia.

Agyneta nigra (Oi, 1960)*

MATERIAL. 1 ♂, B.R., Bureya River Valley, 1 km downstream of confluence of P. and L. Bureya River, 16–21.VI.2004, leg. L. Trilikauskas.

DISTRIBUTION. East Siberia, Sakhalin, Lake Baikal region; Amurskaya, Khabarovsk and Maritime provinces. Known also from Mongolia, China, Japan and Korea.

Agyneta olivacea (Emerton, 1882)

2004 *Agyneta olivacea* — Tanasevitch & Trilikauskas: 78.

2005 *Agyneta olivacea* — Trilikauskas: 212.

MATERIAL. 1 ♂, B.R., P. Bureya River Valley, upper reaches of Lednikovoy River, 22.VI.2000, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Allomengea dentisetis (Grube, 1889)

2004 *Allomengea dentisetis* — Tanasevitch & Trilikauskas: 78.

MATERIAL. 3 ♂♂, 2 ♀♀, Khabarovsk Province, Chegdomyn Village, 13–20.VIII.2001, 11.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Anguliphantes dybowskii (O. Pickard-Cambridge, 1873)

2004 *Anguliphantes dybowskii* — Tanasevitch & Trilikauskas: 78.

2005 *Anguliphantes dybowskii* — Trilikauskas: 212.

MATERIAL. 1 ♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas; 1 ♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Anguliphantes karpinskii (O. Pickard-Cambridge, 1873)

2004 *Anguliphantes karpinskii* — Tanasevitch & Trilikauskas: 79.

MATERIAL. 1 ♂, 2 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18.VIII.2004, leg. L. Trilikauskas; 1 ♂, same locality, Chapkhoz River Valley, 300 m upstream of the mouth, 22.VIII.2004, leg. L. Trilikauskas; 1 ♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Bathyphantes eumenis (L. Koch, 1879)

2004 *Bathyphantes simillimus* — Tanasevitch & Trilikauskas: 79.

2005 *Bathyphantes eumenis* — Trilikauskas: 213.

MATERIAL. 4 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18–19.VIII.2004, leg. L. Trilikauskas; 2 ♀♀, B.R., Bureya River Valley, 4 km downstream of confluence of P. & L. Bureya, near mouth of Umalta-Makit River, 06.VIII.2004, leg. D. Logunov; 7 ♀♀, B.R., P. Bureya River Valley, upper reaches of Lednikovoy River, 8–21.VII.2005, leg. L. Trilikauskas; 1 ♀, same locality, near mouth of Bureika River, 21.VIII.2005, leg. L. Trilikauskas; 2 ♀♀, B.R., P. Bureya River Valley, near mouth of Lednikovoy River mouth, 26–28.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Fennoscandia, Central Europe, European tundra. Throughout all Siberia: from the Urals to East Siberia, Cisokhotia, Kamchatka, Commander and Kurile Islands, Sakhalin; through the mountains of South Siberia eastward to Khabarovsk and Maritime provinces. Known from Mongolia and China. Northern Nearctic: from Alaska eastward to Newfoundland.

Bathyphantes gracilis (Blackwall, 1841)*

MATERIAL. 1 ♀, Khabarovsk Province, near the Chegdomyn Village, 08.IX.2002, leg. L. Trilikauskas.

DISTRIBUTION. Through all Europe, including the tundra zone, Turkey, the Caucasus; in Siberia: from the Urals eastward to East Siberia, Kamchatka, Sakhalin; through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk and Maritime provinces; known from China. Northern Nearctic: from Alaska eastward to Newfoundland.

Centromerus pacificus Eskov et Marusik, 1992

MATERIAL. 7 ♂♂, 1 ♀, B.R., Bureya River Valley, 1 km downstream of confluence of P. & L. Bureya, 1–16.VI.2004, leg. L. Trilikauskas.

DISTRIBUTION. East Siberia, Cisokhotia; Mountains of South Siberia: Sayan; Amurskaya, Khabarovsk and Maritime provinces, Sakhalin.

Ceraticelus orientalis Eskov, 1987

2004 *Ceraticelus orientalis* — Tanasevitch & Trilikauskas: 80.

MATERIAL. 3 ♂♂, 3 ♀♀, B.R., confluence of P. & L. Bureya, 17.VII.2003, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Ceratinella wideri (Thorell, 1871)

2004 *Ceratinella brevipes* — Tanasevitch & Trilikauskas: 80.

2004 *Ceratinella scabrosa* — Tanasevitch & Trilikauskas: 80.

MATERIAL. 1 ♀, B.R., P. Bureya River Valley, upper reaches of Lednikovoy River, 08.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Widespread in Europe, in Siberia known from the Urals eastward to Chukotka, Cisokhotia; from eastern Kazakhstan through the mountains of South Siberia eastward to Khabarovsk Prov. Recorded in Mongolia and Middle Asia.

Diplocentria bidentata (Emerton, 1882)*

MATERIAL. 2 ♀♀, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. & L. Bureya rivers, 25.VII.2004, 05.VIII.2004, leg. D. Logunov.

DISTRIBUTION. Northern Europe; Siberia: from the Urals to East Siberia, Cisokhotia, Kamchatka, the Kurile Islands, Sakhalin. Through the mountains of South Siberia via the Lake Baikal region eastward to Maritime Prov. Known from Mongolia and China. Widespread in the Nearctic.

Diplocephalus subrostratus (O. Pickard-Cambridge, 1873)

2004 *Diplocephalus subrostratus* — Tanasevitch & Trilikauskas: 80.

MATERIAL. 12 ♂♂, 3 ♀♀, Khabarovsk Province, upper reaches of Bureya River, Umalta – Makit River Valley, 1 km upstream of the mouth, 10.VIII.2004, leg. D. Logunov.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Epibellowia pacifica (Eskov et Marusik, 1992)

2004 *Epibellowia pacifica* — Tanasevitch & Trilikauskas: 80.

MATERIAL. 1 ♂, B.R., L. Bureya River Valley, near mouth of Vankish River, 21.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Erigone atra Blackwall, 1833

2004 *Erigone atra* — Tanasevitch & Trilikauskas: 80.

2005 *Erigone atra* — Trilikauskas: 213.

MATERIAL. 3 ♂♂, Khabarovsk Province, Chegdomyn Village, 11.V.2000, 15.VII.2001, 2–7.VIII.2001, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Erigone capra Simon, 1884*

2004 *Erigone capra* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 1 ♀, B.R., P. Bureya River Valley, near mouth of Lednikovy River, 18–28.VII.2005, leg. L. Trilikauskas 1 ♂, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. & L. Bureya, 22.V–02.VI.2003, leg. A. Tanasevitch.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Erigone dentipalpis (Wider, 1834)

MATERIAL. 1 ♂, B.R., Bureya River Valley, 2.5 km downstream of confluence of P. & L. Bureya, 22.V–02.VI.2003, leg. A. Tanasevitch.

DISTRIBUTION. Throughout the Palaearctic. In the Nearctic known only from Newfoundland.

Floronia bucculenta (Clerck, 1757)

2004 *Floronia bucculenta* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 1 ♀, Khabarovsk Province, Chegdomyn Village, 13.VIII.2000, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Gnathonarium dentatum (Wider, 1834)

2004 *Gnathonarium dentatum* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 2 ♀♀, Khabarovsk Province, Chegdomyn Village, 08.IX.2000, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Gnathonarium taczanowskii (O. Pickard-Cambridge, 1873)

2004 *Gnathonarium taczanowskii* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 8 ♀♀, B.R., P. Bureya River Valley, near mouth of Lednikovy River, 30.VI–18.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Halorates submissus L. Koch, 1879

2004 *Halorates submissus* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 1 ♂, Khabarovsk Province, Chegdomyn Village, 03.V.2001, leg. L. Trilikauskas; 1 ♀, Bureya River Valley, 6 km upstream of Usman River mouth, 28.VI.2001, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Helophora insignis (Blackwall 1841)*

MATERIAL. 5 ♂♂ (subad.), Khabarovsk Province, near the Chegdomyn Village, 09.VIII.2000, leg. L. Trilikauskas.

DISTRIBUTION. Throughout the Holarctic.

Hilaira devitata Eskov, 1987*

MATERIAL. 3 ♀♀, B.R., P. Bureya River Valley, near mouth of Lednikovy River, 10–18.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Middle and East Siberia, southern part of Khabarovsk and Maritime provinces.

Hilaira herniosa (Thorell, 1875)

2004 *Hilaira herniosa* — Tanasevitch & Trilikauskas: 81.

MATERIAL. 2 ♂♂, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Holminaria sibirica Eskov, 1991

2004 *Holminaria sibirica* — Tanasevitch & Trilikauskas: 82.

MATERIAL. 1 ♂, 1 ♀, Khabarovsk Province, upper reaches of Bureya River, Umalta-Makit River Valley, 1 km upstream of the mouth, 10.VIII.2004, leg. D. Logunov; 1 ♀, same locality, Chegdomyn Village, 22.V.2005, leg. L. Trilikauskas; 7 ♀♀, B.R., P. Bureya River Valley, upper reaches of Lednikovy River, 10.VII–07.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Hypomma bituberculatum (Wider, 1834)

2000 *Hypomma bituberculatum* — Trilikauskas: 217.

2005 *Hypomma bituberculatum* — Trilikauskas: 213.

MATERIAL. 1 ♀, Khabarovsk Province, Verchnebureinskiy Distr., near mouth of Niman River, 21.VI.2003, leg. L. Trilikauskas.

DISTRIBUTION. Throughout the Palaearctic.

Hylyphantes nigrinus (Simon, 1881)*

MATERIAL. 1 ♀, Khabarovsk Province, near the Chegdomyn Village, 13.VIII.2000, leg. L. Trilikauskas.

DISTRIBUTION. Widespread in Europe, the Caucasus, the Urals. Eastern Kazakhstan, throughout the mountains of South Siberia via the Lake Baikal region to Khabarovsk and Maritime provinces, the Kurile Islands, Sakhalin. Known from China.

Hypselistes jacksoni (O. Pickard-Cambridge, 1902)

2004 *Hypselistes jacksoni* — Tanasevitch & Trilikauskas: 82.

MATERIAL. 1 ♀, Khabarovsk Province, Verchnebureinskiy Distr., near mouth of Niman River, 20.VI.2003, leg. L. Trilikauskas; 2 ♂♂, 2 ♀♀, B.R., L. Bureya River Valley, near mouth of Vankish River, 21.VIII.2004, leg. L. Trilikauskas; 3 ♂♂, B.R., L. Bureya River basin, Chapkhoz River Valley, 200 m upstream of its mouth, 22.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Hypselistes semiflavus (L. Koch, 1879)

2004 *Hypselistes semiflavus* — Tanasevitch & Trilikauskas: 82.
MATERIAL. 1 ♀, Bureya River Valley, near mouth of Niman River, 20.VI.2003, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Incestophantes incestoides (Tanasevitch et Eskov, 1987)*

MATERIAL. 1 ♀, B.R., P. Bureya River Valley, 1.5 km upstream of the mouth, 20.VII.2004, leg. D. Logunov.

DISTRIBUTION. The Urals, East Siberia, Cisokhotia, Khabarovsk Prov.

Incestophantes laricetorum (Tanasevitch et Eskov, 1987)

2004 *Incestophantes laricetorum* — Tanasevitch & Trilikauskas: 82.

2005 *Incestophantes laricetorum* — Trilikauskas: 213.
MATERIAL. 1 ♂, 1 ♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Incestophantes obtusus Tanasevitch, 1996

2004 *Incestophantes obtusus* — Tanasevitch & Trilikauskas: 82.
2005 *Incestophantes obtusus* — Trilikauskas: 213.

MATERIAL. 1 ♀, B.R., L. Bureya River Valley, near mouth of Vankish River, 22.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Linyphia triangularis (Clerck, 1757)

2005 *Linyphia triangularis* — Trilikauskas: 213.
MATERIAL. 1 ♂, Khabarovsk Province, Chegdomyn Vilage, 13.VIII.2000, leg. L. Trilikauskas.

DISTRIBUTION. Throughout the Palearctic, introduced in USA.

Macrargus multesimus (O. Pickard-Cambridge, 1875)*

MATERIAL. 1 ♀, B.R., Bureya River Valley, 1 km downstream of confluence of P. and L. Bureya River, 12–17.VII.2004, leg. D. Logunov.

DISTRIBUTION. Europe include tundra zona; Siberia: from the Urals eastward to Chukotka, Cisokhotia, Kamchatka, Sakhalin. Through the mountains of South Siberia eastward to Khabarovsk and Maritime provinces. Known from Mongolia and China.

Maro borealis Eskov, 1991

2004 *Maro borealis* — Tanasevitch & Trilikauskas: 83.
MATERIAL. 1 ♂, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18.VIII.2004, leg. L.

Trilikauskas; 4 ♀♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VIII.2005, leg. L. Trilikauskas

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Maro bureensis Tanasevitch, 2006

2004 *Maro* sp. — Tanasevitch & Trilikauskas: 83.

2006 *Maro bureensis* — Tanasevitch: 262.

MATERIAL. 2 ♀♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Southern part of Khabarovsk Province.

Maro khabarum Tanasevitch, 2006

2006 *Maro khabarum* — Tanasevitch: 262.

MATERIAL. 1 ♂, L. Bureya River basin, Chapkhoz River Valley, 22.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. Amurskaya and Khabarovsk provinces.

Maro pansibiricus Tanasevitch, 2006

2006 *Maro pansibiricus* — Tanasevitch: 260.

MATERIAL. 1 ♂, 1 ♀, B.R., L. Bureya River Valley, near mouth of Vankish River, 21.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. West, Middle and East Siberia (except Chukotka), through the mountains of South Siberia eastward to Khabarovsk and Maritime provinces, Sakhalin.

Maso sundevalli (Westring, 1851)

2004 *Maso sundevalli* — Tanasevitch & Trilikauskas: 83.

MATERIAL. 1 ♀, L. Bureya River basin, Chapkhoz River Valley, 22.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. All over the Holarctic.

Mecynargus monticola (Holm, 1943)*

MATERIAL. 1 ♀, B.R., Bureya River Valley, 2.5 km downstream of confluence of P. and L. Bureya River, 16–21.VII.2002, leg. L. Trilikauskas; 2 ♀♀, same locality, 23.V.2003, 31.V.2003, leg. A. Tanasevitch; 1 ♂, same locality, 05.VIII.2004, leg. D. Logunov; 1 ♂, B.R., P. Bureya River Valley, near mouth of Lednikovoy River, 9–18.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Northern Europe; Siberia: from the Urals to Chukotka, Cisokhotia. Through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk Prov. Known from Mongolia. In the Nearctic known from Canada: British Columbia, Yukon, Northern Territories.

Mecynargus pinipumilis Eskov, 1988*

MATERIAL. 1 ♂, 2 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas; 3 ♂♂, 2 ♀♀, same locality, near mouth of Vankish River, 22.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. East Siberia, Cisokhotia, Khabarovsk Prov.

Metopobactrus prominulus (O. Pickard-Cambridge, 1872)*

MATERIAL. 1 ♀, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. and L. Bureya River, 25.VII.2004, leg. D. Logunov.

DISTRIBUTION. Throughout the Holarctic.

Micrargus herbigradus (Blackwall, 1854)

2004 *Micrargus herbigradus* — Tanasevitch & Trilikauskas: 83.
 MATERIAL. 1 ♂, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VIII.2005, leg. L. Trilikauskas.
 DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Microlinyphia pusilla (Sundevall, 1830)

2004 *Microlinyphia pusilla* — Tanasevitch & Trilikauskas: 83.
 MATERIAL. 1 ♀, B.R., P. Bureya River Valley, 3–4 km upstream of the mouth, 10–14.VII.1999, leg. L. Trilikauskas.
 DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Microneta viaria (Blackwall, 1841)

2004 *Microneta viaria* — Tanasevitch & Trilikauskas: 83.
 MATERIAL. 1 ♂, B.R., P. Bureya River Valley, 300 m upstream of its mouth, 2–7.VII.2004, leg. L. Trilikauskas; 3 ♂♂, Khabarovsk Province, Verchnebureinskiy Distr., Umalta-Makit River Valley, 1 km upstream of its mouth, 10.VIII.2004, leg. L. Trilikauskas.
 DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Minicia marginella (Wider, 1834)*

MATERIAL. 1 ♀, Khabarovsk Province, Bureya River Valley, near mouth of Umalta-Makit River, 03.VIII.2004, leg. D. Logunov.

DISTRIBUTION. Europe, eastern Kazakhstan, Middle Asia, the Urals; West and East Siberia, Cisokhotia, Khabarovsk Prov.; in South Siberia registered from Tuva and Sayan.

Minyrioloides trifrons (O. Pickard-Cambridge, 1863)*

MATERIAL. 1 ♂, 1 ♀, B.R., P. Bureya River Valley, Bureika River basin, Siuriuktak River Valley, 300 m upstream of its mouth, 30.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Throughout the Holarctic.

Mughiphantes aculifer (Tanasevitch, 1988)

2004 *Mughiphantes aculifer* — Tanasevitch & Trilikauskas: 83.
 2005 *Mughiphantes aculifer* — Trilikauskas: 213.
 MATERIAL. 1 ♂, 3 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18–19.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Mughiphantes sachalinensis (Tanasevitch, 1988)

2004 *Mughiphantes sachalinensis* — Tanasevitch & Trilikauskas: 83.

MATERIAL. 1 ♂, B.R., Bureya River Valley, 2.5 km downstream of confluence of P. and L. Bureya River, 06.IX.2003, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Mughiphantes taczanowskii (O. Pickard-Cambridge, 1873)

2004 *Mughiphantes taczanowskii* — Tanasevitch & Trilikauskas: 83.

MATERIAL. 6 ♀♀, B.R., L. Bureya River Valley, near mouth of Vankish River, 22.VIII.2004, leg. L. Trilikauskas; 1 ♀, same locality, Chapkhoz River Valley, 200 m upstream of its mouth,

22.VIII.2004, leg. L. Trilikauskas; 1 ♀, B.R., P. Bureya River Valley, Bureika River basin, Siuriuktak River Valley, 300 m upstream mouth, 30.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Nematogmus sanguinolentus (Walckenaer, 1841)*

2004 *Nematogmus sanguinolentus* — Tanasevitch & Trilikauskas: 83.

MATERIAL. 1 ♀, Khabarovsk Province, Chegdomyn Village, 31.V.2005, leg. L. Trilikauskas; 1 ♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Nenilinium asiaticum Eskov, 1988

2004 *Nenilinium luteolum* — Tanasevitch & Trilikauskas: 30.
 MATERIAL. 12 ♂♂, 27 ♀♀, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. & L. Bureya, 22–27.V.2003, leg. A. Tanasevitch; 1 ♀, same locality, 24.VII.2004, leg. D. Logunov.

DISTRIBUTION. Middle and East Siberia; through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk Prov., Sakhalin.

Neriene clathrata (Sundevall, 1830)

2004 *Neriene clathrata* — Tanasevitch & Trilikauskas: 84.

MATERIAL. 1 ♀, Khabarovsk Province, Chegdomyn Village, 02.VI.2000, leg. L. Trilikauskas; 1 ♂, B.R., L. Bureya River Valley, Chapkhoz River Valley, 5–6 km upstream of its mouth, 11–12.VI.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Neriene oidedicata van Helsdingen, 1969*

MATERIAL. 1 ♀, Khabarovsk Province, Chegdomyn Village, 03.IX.2001, leg. L. Trilikauskas; 3 ♂♂, 3 ♀♀, Bureya River Valley, near mouth of Niman River, 21–22.VI.2003, leg. L. Trilikauskas.

DISTRIBUTION. Known from China, Japan and Korea

Oreonetides helsdingeni Eskov, 1984

2004 *Oreonetides helsdingeni* — Tanasevitch & Trilikauskas: 84.

2005 *Oreonetides helsdingeni* — Trilikauskas: 214.

MATERIAL. 1 ♂, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Oreonetides kolymensis Eskov, 1991*

MATERIAL. 1 ♂, B.R., Bureya River Valley, 1 km downstream of confluence of P. and L. Bureya River, 16–21.VI.2004, leg. L. Trilikauskas.

DISTRIBUTION. East Siberia, Chukotka, Cisokhotia, Khabarovsk Prov.

Praestigia pini (Holm, 1950) *

MATERIAL. 1 ♂, B.R., Bureya River Valley, 2.5 km downstream of confluence of P. and L. Bureya River, 6–11.VI.2004, leg. L. Trilikauskas.

DISTRIBUTION. Northern Europe; in Siberia: from the Urals eastward to East Siberia; through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk Prov.; Sakhalin. Known also from Mongolia.

Savignia basarukini Eskov, 1988*

MATERIAL. 2 ♂♂, B.R., P. Bureya River Valley, 300 m upstream of the mouth, 01.VI.2003, leg. A. Tanasevitch.

DISTRIBUTION. Amurskaya and Khabarovsk provinces, Sakhalin.

Savignia bureensis Tanasevitch et Trilikauskas, 2006

2006 *Savignia bureensis* — Tanasevitch et Trilikauskas: 269.

COMMENTS. This species has been recently described from the Bureinsky Reserve by Tanasevitch & Trilikauskas [2006]: upper reaches of Bureya River, mouth of Levyi Ussomakh Stream, *Larix* forest, in lichen.

DISTRIBUTION. Amurskaya and Khabarovsk provinces.

Silometopoides sibiricus (Eskov, 1989)*

MATERIAL. 2 ♂♂, 4 ♀♀, B.R., Bureya River Valley, 2.5 km downstream of confluence of P. and L. Bureya River, 25.VII.2004, leg. D. Logunov.

DISTRIBUTION. The Urals, East Siberia, Sayan (South Siberia), Khabarovsk Prov.

Styloctetor romanus (O. Pickard-Cambridge, 1872)*

MATERIAL. 2 ♂♂, Khabarovsk Province, Bureya River Valley, near mouth of Umalta-Makit River, 03.VIII.2004, leg. D. Logunov.

DISTRIBUTION. Throughout the Palaearctic.

Theoneta aterrima (Eskov et Marusik, 1991)*

MATERIAL. 1 ♂, B.R., P. Bureya River Valley, upper reaches of Lednikovoy River, 11–17.VI.2000, leg. L. Trilikauskas; 2 ♂♂, upper reaches of the Bureya River, Umalta – Makit River Valley, 10.VIII.2004, leg. D. Logunov; 2 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18.VIII.2004, leg. L. Trilikauskas; 1 ♂, 2 ♀♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VII.2005, 29.VII–02.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Amurskaya and Khabarovsk provinces.

Theoneta saaristoi (Eskov et Marusik, 1991)

2004 *Theoneta saaristoi* — Tanasevitch & Trilikauskas, *Arthropoda Selecta*, 13 (1–2): 85.

MATERIAL. 1 ♂, 6 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Tibioplus diversus (L. Koch, 1879)*

MATERIAL. 1 ♀, B.R., P. Bureya River Valley, near mouth of Bureika River, 21.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Throughout the Holarctic.

Tmeticus nigriceps (Kulczyński, 1916)*

MATERIAL. 1 ♀, B.R., near mouth of P. Bureya River, 30.VI.2002, leg. L. Trilikauskas.

DISTRIBUTION. Siberia: from the Urals eastward to Chukotka, Khabarovsk Prov.

Tmeticus tolli Kulczyński, 1908*

MATERIAL. 1 ♂, B.R., L. Bureya River Valley, near mouth of Vankish River, 21.VIII.2004, leg. L. Trilikauskas; 1 ♂, same locality, Chapkhoz River Valley, 22.VIII.2004, leg. L. Trilikauskas; 5 ♂♂, B.R., P. Bureya River Valley, Bureika River basin, Siuriuktak River Valley, 300 m upstream of its mouth, 30.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. Siberia: from Yenisey River to Chukotka, Cisokhotia; Kamchatka, Sakhalin; Amurskaya and Khabarovsk provinces.

Trematocephalus cristatus (Wider, 1834)*

MATERIAL. 2 ♀♀, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. and L. Bureya River, 2–5.VIII.2004, leg. D. Logunov; 2 ♀♀, B.R., P. Bureya River Valley, 300 m upstream of the mouth, 06.VIII.2004, leg. D. Logunov & L. Trilikauskas.

DISTRIBUTION. Widespread in Europe, known from the Caucasus and the Urals. In Siberia: West Siberia; through the mountains of South Siberia via the Lake Baikal region eastward to Amurskaya, Khabarovsk and Maritime provinces. Known also from China.

Ummeliata sibirica (Eskov, 1980)*

MATERIAL. 1 ♀, Khabarovsk Province, Chegdomyn Village, IX.2002, leg. L. Trilikauskas.

DISTRIBUTION. Middle Siberia; through the mountains of South Siberia via the Lake Baikal region eastward to Khabarovsk and Maritime provinces.

Viktorium putoranicum Eskov, 1988*

MATERIAL. 1 ♂, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. The Urals, Middle and East Siberia; Tuva, Sayan (South Siberia); Khabarovsk Prov., Sakhalin.

Wabasso tungusicus Eskov, 1988*

MATERIAL. 1 ♀, B.R., Bureya River Valley, 3.5 km downstream of confluence of P. and L. Bureya River, 25.V.2003, leg. A. Tanasevitch.

DISTRIBUTION. Middle Siberia, Cisokhotia, Amurskaya and Khabarovsk provinces.

Walckenaeria auranticeps (Emerton, 1882)*

MATERIAL. 1 ♂, Khabarovsk Province, Chegdomyn Village, 01.VI.2005, leg. L. Trilikauskas

DISTRIBUTION. East Siberia, Cisokhotia; in South Siberia known from Tuva and Sayan, Amurskaya and Khabarovsk provinces. Widespread in the Nearctic.

Walckenaeria karpinskii (O. Pickard-Cambridge, 1873)

2004 *Walckenaeria karpinskii* — Tanasevitch & Trilikauskas: 86.

2005 *Walckenaeria karpinskii* — Trilikauskas: 214.

MATERIAL. 1 ♂, 2 ♀♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 19.VIII.2004, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Walckenaeria korobeinikovi Esyunin Efimik, 1996

2004 *Walckenaeria korobeinikovi* — Tanasevitch & Trilikauskas: 86.

2005 *Walckenaeria korobeinikovi* — Trilikauskas: 214.

MATERIAL. 1 ♀, B.R., L. Bureya River Valley, 3 km upstream of Kuraigagna River mouth, 18–19.VIII.2004, leg. L. Trilikauskas; 1 ♀, B.R., P. Bureya River Valley, near mouth of Lednikov River, 18–25.VII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

Zornella cultrigera (L. Koch, 1879)

2004 *Zornella cultrigera* — Tanasevitch & Trilikauskas: 86.

2005 *Zornella cultrigera* — Trilikauskas: 214.

MATERIAL. 1 ♀, B.R., P. Bureya River Valley, upper reaches of Lednikov River, 8–21.VII.2005, leg. L. Trilikauskas; 1 ♀, same locality, near mouth of Lednikov River, 9–18.VII.2005, leg. L. Trilikauskas; 3 ♀♀, same locality, near mouth of Bureika River, 29.VII–02.VIII.2005, leg. L. Trilikauskas.

DISTRIBUTION. See Tanasevitch & Trilikauskas [2004].

ACKNOWLEDGEMENTS. This work has been supported in part by the Russian Foundation for the Basic Research (RFFI). Project 02-04-49742.

References

- Tanasevitch A.V. 2006. New or little-known *Maro* O.P.-Cambridge from Siberia and the Russian Far East (Aranei: Linyphiidae: Micronetinae) // *Arthropoda Selecta*. Vol.14. No.3. P.259–268.
- Tanasevitch A.V., Trilikauskas L.A. 2004. Check-list of the linyphiid spiders of the Bureinsky State Nature Reserve and adjacent territories, Russian Far East (Aranei: Linyphiidae) // *Arthropoda Selecta*. Vol.13. No.1–2. P.77–86.
- Tanasevitch A. V., Trilikauskas L.A. 2006. A new species of the genus *Savignia* Blackwall from the Khabarovsk Province, Russian Far East (Aranei: Linyphiidae: Erigoninae) // *Arthropoda Selecta*. Vol.14. No.3. P.269–270.
- Trilikauskas L.A. 2000. Notes on the spider fauna (Aranei) of the upper reaches of Bureya River (Khabarovsk Province) // *Arthropoda Selecta*. Vol.9. No.3. P.215–220.
- Trilikauskas L.A. 2005. Spiders of the larch forests of the Bureinsky reserve and contiguous territories: annotated list of species // *Issledovanie i konstruirovanie landshaftov Dalnego Vostoka i Sibiri*. Ed. Urusov V.M. Vladivostok. Dalnauka. No.6. P.207–217 [in Russian].